



## **Shoulder Replacement Post-Operative Instructions**

### **Initial Instructions**

- Ice along with pain medicine are your keys to initial pain management.
- After you arrive home, rest the remainder of the day. You may be tired or nauseated from the anesthesia and/or pain medication.
- You will be in a sling after surgery. You will typically be in the sling for **6 weeks** post-operatively.

### **Dressing Change**

- Leave the dressing in place for 3-5 days following surgery.
- The dressing is waterproof and you may shower without covering the dressing. If the gauze under the dressing happens to get wet or dirty, just remove it and replace it with a clean dressing.
- After 3-5 days, remove the dressing making sure to leave the steri-strips on the incision in place.
- It is extremely important to keep the surgical site clean and dry at all times to help prevent infection.
- Signs of infection include redness, swelling, increased pain, warmth, fever (above 101.4 degrees F) and drainage from the wound. If you note any of these signs, please call the office immediately at (214) 265-3200.

### **Bathing**

- You may shower **immediately after your surgery** with the waterproof dressing in place. After 3-5 days and your dressing is removed, you may get your incision wet as long as the wounds are dry and not draining. Let the water run down over the incision but do not rub or scrub them.
- If the wounds are draining, then wait until they are dry before getting them wet.
- **Do not soak or submerge** the surgical site until the wounds are completely healed (typically 4-6 weeks after surgery).

### **Activity**

- Keep sling in place at all times as directed.
- OK to remove sling with minimal movement about arm for dressing, hygiene, and prescribed exercises.
- May return to **sedentary work ONLY** 3-4 days after surgery, if pain is tolerable
- **OK to use computer.** Use your other hand to place your operated extremity on the desk or keyboard and allow it to fully rest there without using your own muscles to keep it in position.

### **Physical Therapy**

- **You will be discharged with a physical therapy prescription and a physical therapy protocol.** The protocol is a description for you and your physical therapist that details all of your restrictions and activities through your entire recovery. Please be sure to give this to your physical therapist.
- **Please arrange your post-operative physical therapy appointment.** Our office can help you with this if needed. You may start as soon as the day following your surgery, but most prefer the

week after surgery. They will guide you in exercises to perform while in the clinic and at home.

## Sling/Immobilizer

- You will be put in a sling after surgery.
- **Sling should be worn at all times (except for hygiene and exercises)**
- Keep your elbow position against the pillow and even with your side or in front of this position to minimize stress on the repair.
- You will typically wear the sling for **6 weeks post-operatively**, unless instructed otherwise.

## Ted Stockings

- These compressive stockings/socks are to be worn for **2 weeks** after surgery
- They aid in the control of lower extremity swelling and decrease the chance of blood clot formation
- You may remove for hygiene, but should be worn at all other times

## Ice Therapy

- Icing is very important to decrease swelling and pain and improve mobility.
- You will be provided with a cryocuff.
- Keep cold therapy cuff on at all times for the first 24 hours after surgery.
- Refill with cold water every hour while awake.
- After 24 hours, continue to use the cuff 3-4 times a day, 15-20 minutes each time.
- Be sure to keep clothing or a towel between ice and skin, as placing the ice directly on skin may cause frostbite. **DO NOT PLACE DIRECTLY ON SKIN.**
- If a cryocuff is not provided, please use ice for 20 minutes each hour over surgical site. **DO NOT PLACE DIRECTLY ON SKIN.**

## Medications

- **If placed, nerve blocks and pain medication injected into the wound will wear off from 12-72 hours after surgery.**
- You will be given a narcotic pain medication. This should be taken as needed every 4-6 hours for pain. Take with a little food.
- Narcotic pain medications **will not alleviate all of your discomfort**
- **Ice and elevation** will aide in additional pain control
- Take medications as prescribed
- **Do not mix narcotics with alcohol**
- Common side effects of narcotic pain medications include
  - Nausea- Take medication with food. If prescribed you may use anti-nausea medication. If nausea persists, call the office to consider having the medication changed.
  - Drowsiness
  - Constipation - We recommend purchasing an over-the-counter stool softener (such as colace) to take as directed while taking narcotics to prevent constipation. You will be given a prescription for this.
- **Do not drive a vehicle or operate heavy machinery while on narcotic pain medications**
- Narcotic medications **cannot** be called into the pharmacy. They will require a hand-written prescription from the office. Please monitor the amount of pain medications you have remaining in order to not run out
- **Narcotics and pain medications will not be refilled on weekends or after hours.**
- You will likely be given a prescription for **aspirin** to take post-operatively. This is typically taken only **once a day** for 3 weeks post-operatively in combination with your compression socks **to help prevent blood clots**. The dosing and duration may change on an individual basis, but it should be reflected in the prescription that you are given.
- We also recommend taking an anti-inflammatory if you are able to tolerate them, such as Motrin or Naproxen, unless you are taking blood thinners or instructed not to take the anti-inflammatories

by your doctor. Take as directed. Take with food.

## Post-Op Office Appointment

- If the appointment was not arranged prior to surgery, please contact the office at (214) 265-3200.
- **Your first post-operative visit should be 10-14 days after surgery**
- Depending on the procedure plan for follow-up visits at **6 weeks, 3 months, and 6 months** after surgery.

## Emergencies or Concerns

Contact us at **(214) 265-3200** or go to the nearest emergency room **if you have any of the following:**

- Pain worsens or does not improve after pain medication is taken.
- Any abrupt change in the nature or severity of pain
- Redness or swelling in your thigh or calf that will not decrease.
- Redness around incisions
- Continuous drainage or bleeding from the incision (occasional drainage is expected)
- Difficulties moving your fingers or **extreme pain when moving your fingers**
- Numbness or tingling that does not go away or is getting worse
- Cold hand or color change in extremity (Please be aware that the skin prep in surgery may have an orange, brown, or blue tint)
- Develop a **fever greater than 101.5 degrees F.**
- Chest pain, shortness of breath, intractable nausea/vomiting, or anything else concerning
- If you have an emergency after office hours or on the weekend, call (214) 265-3200 to be connected to our pager service. They will contact Dr. Raynor or one of his colleagues. **Do not** call the hospital or surgery center.
- **If you have an emergency that requires immediate attention, proceed to the nearest emergency room.**

Dr. Raynor strives to take **exceptional** care of his patients. If there are any concerns, difficulties, or problems, please let him know. He wants to work with you to achieve the best possible outcomes. This is best obtained with an open line of communication.

***Wishing you a quick and successful recovery.***

**M. Brett Raynor, MD**

Orthopaedic Surgery and Sports Medicine Specialist  
Texas Orthopaedic Associates, L.L.P.

[www.brettraynormd.com](http://www.brettraynormd.com)

[www.txorthopaedic.com](http://www.txorthopaedic.com)

